

**K/2 metals**

Na <sub>2</sub> EDTA · 2H <sub>2</sub> O	3.722 g
CuSO <sub>4</sub> · 5H <sub>2</sub> O	0.25 mg
Na <sub>2</sub> MoO <sub>4</sub> · 2H <sub>2</sub> O	0.726 mg
ZnSO <sub>4</sub> · 7H <sub>2</sub> O	2.3 mg
CoSO <sub>4</sub> · 7H <sub>2</sub> O	1.406 mg
MnCl <sub>2</sub> · 4H <sub>2</sub> O	17.81 mg
H <sub>2</sub> SeO <sub>3</sub>	0.129 mg
NiCl <sub>2</sub> · 6H <sub>2</sub> O <sup>1)</sup>	0.149 mg
Distilled water	100 mL

1) In the NIES-Collection, 0.149 mg . NiCl<sub>2</sub> · 6H<sub>2</sub>O is replaced by 0.812 mg NiCl<sub>2</sub>.

**Reference**

Keller, M. D., Guillard, R .R. L. 1985. Factors significant to marine diatom culture. pp. 113-6. *In* Anderson, D. M., White, A. W., Baden, D. G. (eds.) *Toxic Dinoflagellates*. Elsevier, New York.

Keller, M. D., Selvin, R. C., Claus, W., Guillard, R. R. L. 1987. Media for the culture of oceanic ultraphytoplankton. *J. Phycol.*, 23, 633-638.